PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:
A61K 31/44, 9/08

A1
(11) International Publication Number: WO 94/02141
(43) International Publication Date: 3 February 1994 (03.02.94)

(21) International Application Number: PCT/JP93/00998 (74) Agent: TAKASHIMA, Hajime; Yuki Building, 3-9, Hira-

JP

15 July 1993 (15.07.93)

(71) Applicants (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP). YOSHITOMI PHARMACEUTICAL INDUSTRIES, LTD. [JP/JP]; 6-9, Hiranomachi 2-chome, Chuo-ku,

28 July 1992 (28.07.92)

Osaka-shi, Osaka 541 (JP). (72) Inventors; and

(22) International Filing Date:

(30) Priority data: 4/201203

(75) Inventors/Applicants (for US only): NAKANISHI, Shigeo [JP/JP]; 12-5, Shimokidacho, Neyagawa-shi, Osaka 572 (JP). TOMINAGA, Tetsuo [JP/JP]; B-806, Kasugaoka-Abankonfoto, 136-3, Kasugaoka 2-chome, Itami-shi, Hyogo 664 (JP). YAMANATA, Iwao [JP/JP]; 5-6-12, Kami-minami, Hirano-ku, Osaka-shi, Osaka 547 (JP). HIGO, Takashi [JP/JP]; 2-2-10, Midorigaoka, Ikeda-shi, Osaka 563 (JP). SHIBATA, Toshiyuki [JP/JP]; 774-105, Oaza-Higashihama, Nakatsu-shi, Oita 871 (JP).

(74) Agent: TAKASHIMA, Hajime; Yuki Building, 3-9, Hiranomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).

(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPl patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: INJECTION AND INJECTION KIT CONTAINING OMEPRAZOLE AND ITS ANALOGS

(57) Abstract

An injection comprising a 2-[(2-pyridyl)methylsulfinyl]-benzimidazole compound or a salt thereof having antiulcer activity and an aqueous solvent added with no nonaqueous solvent, wherein the pH of the injection is not less than 9.5 and not more than 11.5, and an injection kit comprising the following (a) and (b), wherein (a) and (b) are adjusted such that the pH upon dissolution of (a) in (b) is 9.5 - 11.5: (a): a lyophilized product of an alkaline aqueous solution of a 2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or salt thereof having antiulcer activity; (b): an aqueous solvent added with no nonaqueous solvent. The injection of the present invention is void of the necessity to lower pH so as to prevent hemolysis and local irritation, and to add a nonaqueous solvent to an aqueous solvent for dissolution so as to prevent concomitant degradation of dissolution property. Accordingly, the injection of the present invention can secure solubility sufficient for formulation into preparation and safety for the human body.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
ΑU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	คบ	Hungary	NZ	New Zealand
BJ	Benin	1E	Ireland	PL	Poland
BR	Brazil	ίŤ	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic	RU	Russian Federation
CF	Central African Republic		of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	ΚZ	Kazakhstan	SI	Slovenia
CI.	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	ĹŸ	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	ÜĀ	Ukraine
DE	Germany	MG	Madagascar	US	United States of America
, DK	Denmark	ML	Mali	UZ	Uzbekistan
ES	Spain	MN	Mongolia	VN	Vict Nam
FI	Finland				

SPECIFICATION

WO 94/02141

INJECTION AND INJECTION KIT CONTAINING OMEPRAZOLE AND ITS ANALOGS

[TECHNICAL FIELD]

The present invention relates to an injection of 2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a salt thereof having antiulcer activity, particularly sodium salt of omeprazole and to an injection kit thereof, which are used in clinical fields.

[BACKGROUND ART]

The 2-[(2-pyridy1)methylsulfinyl]benzimidazole compounds such as omeprazole or lansoprazole are potent antiulcer agents, and are used as pharmaceutical compositions for oral administration. Further, the injections thereof have recently developed.

As an injection of omeprazole, there has been known an injection prepared by dissolving sodium salt of omeprazole in sterilized water, filtering and lyophilizing the solution to give a lyophilized product, and then dissolving the lyophilized product in a mixture of polyethylene glycol 400 for injection, sodium dihydrogenphosphate and sterilized water (Japanese Patent Unexamined Publication No. 167587/1984).

Also, an injection prepared by dissolving a lyophilized product of an alkaline aqueous solution of a 2-[(2-pyridyl)-methylsulfinyl]benzimidazole compound having antiulcer activity such as lansoprazole in a mixture of (a) acid, and (b) at least one of ethanol, propylene glycol and polyethylene glycol

(Japanese Patent Unexamined Publication No. 138213/1990).

In general, the pH of injection is preferably about 4-8, and a pH above 9 has a probability of causing hemolysis and local irritation.

In the case of the 2-[(2-pyridy1)methylsulfiny1]benzimidazole compound or a salt which may be hereinafter referred
to as "benzimidazole compound or salt thereof" represented by
sodium salt of omepazole, it shows a solubility of the level
permitting formulation into preparation, in water in an alkaline
range of pH 9.5 or above, whereas it shows extremely low
solubility in water at a pH of not more than 9, thus rendering
formulation into preparation very difficult.

While the benzimidazole compound or salt thereof is stable in the alkaline range, it poses a problem in that its stability decreases with the lowering pHs.

For this reason, it has been employed in conventional injections of benzimidazole compound or salt thereof such as sodium salt of omeprazole to add an acid such as hydrochloric acid or sodium dihydrogenphosphate to the solution to keep the pH from neutral to weak basic, and to further add a nonaqueous solvent such as polyethylene glycol, ethanol or propylene glycol in order to obtain a certain level of solubility in such pH range.

Yet, these injections pose problems of local irritation and hemolysis caused by the nonaqueous solvent added to the solution for dissolution.

Accordingly, an object of the invention is to provide an injection of benzimidazole compound or salt thereof, particularly sodium salt of omeprazole causing less side-effects such as hemolysis, and less local irritation, which permits easy formulation.

[DISCLOSURE OF THE INVENTION]

As a result of the intensive study conducted by the inventors with the aim of achieving the aforementioned object, it has now been found that a product obtained by lyophilizing an alkaline aqueous solution of benzimidazole compound or salt thereof, and dissolving same in an aqueous solvent added with no nonaqueous solvent scarcely shows hemolytic property and local irritation, notwithstanding the high pH of from 9.5 to 11.5.

Accordingly, the present invention is:

- (1) an injection comprising a 2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a salt thereof having antiulcer
 activity and an aqueous solvent added with no nonaqueous
 solvent, which has a pH of not less than 9.5 and not more than
 11.5,
- (2) an injection kit comprising the following (a) and (b), wherein (a) and (b) are adjusted such that the pH upon dissolution of (a) in (b) is not less than 9.5 and not more than 11.5;
- (a): a lyophilized product of an alkaline aqueous solution of a2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a salt

thereof having antiulcer activity

(b) : an aqueous solvent added with no nonaqueous solvent.

The 2-[(2-pyridyl)methylsulfinyl]benzimidazole compounds having antiulcer activity which are the element constituting the present invention include, for example, the compounds described in Japanese Patent Unexamined Publication No. 62275/1977,

Japanese Patent Unexamined Publication No. 1417/1979, Japanese Patent Unexamined Publication No. 53406/1982, Japanese Patent Unexamined Publication No. 135881/1983, Japanese Patent Unexamined Publication No. 192880/1983, Japanese Patent Unexamined Publication No. 192880/1983, Japanese Patent Unexamined Publication No. 181277/1984 or Japanese Patent Unexamined Publication No. 50978/1986, and omeprazole [chemical name: 2-[2-(3,5-dimethyl-4-methoxy)-pyridylmethylsulfinyl]-(5-methoxy)benzimidazole] and lansoprazole [chemical name: 2- {2-[(3-methyl-4-(2,2,2-trifluoroethoxy)]-pyridylmethylsulfinyl} - benzimidazole] are exemplified.

As the salts of said benzimidazole compounds, for example, salts of alkaline metal such as sodium salt or potassium salt or salts of alkaline earth metal such as calcium salt or magnesium salt.

In view of the solubility, it is preferable for the present invention to use the salt of benzimidazole compound.

The injection of the present invention has a pH of not less than 9.5 and not more than 11.5, preferably not less than 10 and not more than 11. Where the pH is less than 9.5, the benzimidazole compound or salt thereof does not sufficiently

dissolve in an aqueous solvent and shows poor stability, while where it is more than 11.5, hemolytic property and local irritation become prominent.

According to the present invention, an injection of the benzimidazole compound or salt thereof can be prepared by dissolving the benzimidazole compound or salt thereof in water for injection, etc. along with a strong alkaline compound such as sodium hydroxide, potassium hydroxide, sodium carbonate or L-arginine, to give an alkaline aqueous solution having a pH adjusted to not less than 10.5 and not more than 12.5, preferably not less than 11 and not more than 12. The alkaline aqueous solution may contain mannitol, glycine, sorbitol, inositol, etc. on demand for better forming of a lyophilized product.

The benzimidazole compound is contained in said alkaline aqueous solution in a proportion of 1-50 mg/ml, preferably 5-40 mg/ml on a free compound basis.

Then, this alkaline aqueous solution is filtered for sterilization, and charged in a vial by 0.5-10 ml. After nitrogen gas displacement to be conducted as necessary, the solution is lyophilized by a method known per se. The lyophilized product thus obtained is the (a): a lyophilized product of an alkaline aqueous solution of the 2-[(2-pyridyl)-methylsulfinyl]benzimidazole compounds or salt thereof having antiulcer activity to be contained in the injection kit of the present invention.

When in use, the injection of the present invention can be produced by dissolving the lyophilized product thus obtained in an aqueous solvent added with no nonaqueous solvent, such as physiological saline, aqueous solution of 5% glucose, or distilled water for injection. Said aqueous solvent corresponds to the (b): an aqueous solvent added with no nonaqueous solvent to be contained in the injection kit of the present invention.

The injection of the present invention can be used, for example, in the form of drip infusion, intravenous injection, intramuscular injection, subcutaneous injection.

The concentration of benzimidazole compound in the injection of the present invention may vary depending upon the administration route, and generally ranges in a proportion of 0.05-10 mg/ml, preferably 0.1-5 mg/ml on a free compound basis.

The benzimidazole compound in the injection of the present invention is administered to an adult at 10-100 mg per day on a free compound basis in a single to three times divided doses, depending upon, for example, the symptoms of the patients.

[BEST MODE FOR CARRYING OUT OF THE INVENTION]

Experimental Example 1

Test preparation

- Preparation obtained in Example 1 to be mentioned later
 Test method
- 1. Hemolysis test

Hemolysis was evaluated by Akaishi method using whole blood

of rabbit. The result is given in Table 1.

2. Local irritation test

Local irritation was evaluated by the comparison of necrotic muscular tissue area at the injection site in 3 rabbits at 2 days after the administration of 1 ml of the test preparation by intramuscular injection, with that in the rabbits administered with 1 ml of physiological saline or 1 ml of a 1.7% acetic acid solution, respectively by intramuscular injection.

The results are summarized in Table 2.

Test results

Table 1

Test preparation	рН	Hemolysis
Ex. 1	10.5	not observed

Table 2

Test preparation	рН	Necrotic area (mm²)
Ex. 1	10.5	63
1.7% acetic acid solution (positive comparison solution)	_	398
physiological saline (negative comparison solution)	_	31

(average of 3 rabbits)

The preparation of the present invention is desirable as an injection, since it does not cause hemolysis at all despite the high pH, and causes less local irritation.

Example 1

1N Sodium hydroxide (2.3 ml) is added to 21.3 g of sodium salt of omeprazole (20 g as omeprazole), and water for injection is added thereto to adjust the pH to 11.5 and the total amount to 1 kg.

After filtration for sterilization, this alkaline aqueous solution is charged in 10 ml vials by 2 g. A rubber plug is half driven in, and nitrogen displacement is performed. Lyophilization by a conventional method and dissolution of the lyophilized product obtained in 10 ml of physiological saline give an omeprazole injection [4 mg (free compound)/ml].

[INDUSTRIAL APPLICABILITY]

The injection of the present invention is void of the necessity to lower pH so as to prevent hemolysis and local irritation, and to add a nonaqueous solvent such as polyethylene glycol to an aqueous solvent for dissolution so as to prevent concomitant degradation of dissolution property. As a result, irritation and hemolysis caused by the nonaqueous solvent can be avoided. Accordingly, the injection of the present invention can secure solubility sufficient for formulation into preparation and safety for the human body.

CLAIMS

- 1. An injection comprising a 2-[(2-pyridyl)methylsulfinyl]-benzimidazole compound or a salt thereof having antiulcer activity and an aqueous solvent added with no nonaqueous solvent, wherein the pH of the injection is not less than 9.5 and not more than 11.5.
- 2. The injection of Claim 1, prepared by dissolving a lyophilized product of an alkaline aqueous solution of the 2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a salt thereof having antiulcer activity in the aqueous solvent added with no nonaqueous solvent.
- 3. An injection kit comprising the following (a) and (b), wherein (a) and (b) are adjusted such that the pH upon dissolution of (a) in (b) is not less than 9.5 and not more than 11.5:
- (a): a lyophilized product of an alkaline aqueous solution of a2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a saltthereof having antiulcer activity
- (b) : an aqueous solvent added with no nonaqueous solvent.
- 4. The injection of Claim 1 or 2, wherein the 2-[(2-pyridyl)methylsulfinyl]benzimidazole compound or a salt thereof is sodium salt of omeprazole.
- 5. The injection kit of Claim 3, wherein the 2-[(2-pyridyl)-methylsulfinyl]benzimidazole compound or the salt thereof is sodium salt of omeprazole.

International Application No

_	to International Patent . 5 A61K31/4	t Classification (IPC) or to both National Cl. 4; A61K9/08	assification and IPC	
	• • • • • • • • • • • • • • • • • • • •			
II. FIELDS	S SEARCHED	Minimum Docume	neation Convolved?	
Classifica	tion System		Classification Symbols	
	-		Acontrol o jacon	
Int.Cl	. 5	A61K		
		Documentation Searched other to the Extent that such Documents a		
III. DOCU		ED TO BE RELEVANT ⁹		
Category °	Citation of D	ocument, ¹¹ with indication, where appropria	ite, of the relevant passages 12	Relevant to Claim
A	16 Augus see cla	ims e 7, line 50 - line 52		1-5
A	7 Novem cited in see cla see page see page	e 6, line 6 - line 15 e 7, line 31 - line 37 e 8, line 1 - line 8	ÄSSLE)	1-5
A	28 Febru	356 143 (TAKEDA) uary 1990 n the application ims		1-5
"A" do co "E" eau fil "L" do co wh cit "O" do ot "P" do	onsidered to be of particular document but publing date to establish the batter which may through the batter means occument referring to an their means occument published prior ter than the priority dat	neral state of the art which is not ular relevance lished on or after the international ow doubts on priority claim(s) or the publication date of another eason (as specified) oral disclosure, use, exhibition or to the international filing date but	"T" later document published after the interns or priority date and not in conflict with the cited to understand the principle or theor invention "X" document of particular relevance; the clai cannot be considered novel or cannot be involve an inventive step "Y" document of particular relevance; the clai cannot be considered to involve an invent document is combined with one or more of ments, such combination being obvious to in the art. "&" document member of the same patent fan	he application but y underlying the imed invention considered to imed invention tive step when the other such docu- o a person skilled
	IFICATION Actual Completion of	the International Search	Date of Mailing of this International Sea	rch Report
	29 SEPTEM		0 7, 10, 93	
Total color	al Searching Authority		Signature of Authorized Officer	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

JP 9300998 SA 76470

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

29/0 29/09/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0382489	16-08-90	CA-A- 2009741 JP-A- 3173817 US-A- 5013743	10-08-90 29-07-91 07-05-91
EP-A-0124495	07-11-84	AU-B- 563842 AU-A- 2525784 CA-A- 1264751 GB-A,B 2137616 JP-C- 1651336 JP-B- 3013233 JP-A- 59167587 SU-A- 1314953 US-A- 4738974	23-07-87 06-09-84 23-01-90 10-10-84 30-03-92 22-02-91 21-09-84 30-05-87 19-04-88
EP-A-0356143	28-02-90	JP-A- 2138213 US-A- 5223515	28-05-90 29-06-93